
1. シミ関連英語原著論文

- 1-1. Imokawa G., Mishima Y.: Loss of melanogenic properties in tyrosinases induced by glycosylation inhibitors within malignant melanoma cells. *Cancer Res.* 42: 1994- 2002, 1982.
- 1-2. Mishima Y., Imokawa G.: Selective aberration and pigment loss in melanosomes of malignant melanoma cells in vitro by glycosylation inhibitors: Premelanosomes as glycoprotein. *J. Invest. Dermatol.* 81: 106-114, 1983.
- 1-3. Imokawa G., Mishima Y.: Functional analysis of tyrosinase isozymes of cultured melanoma cells during the recovery period following interrupted melanogenesis induced by glycosylation inhibitors *J. Invest. Dermatol.* 83:196-201, 1984.
- 1-4. Imokawa G., Mishima Y.: Analysis of tyrosinase as asparagine-linked oligosaccharides by Concanavalin-A lectin chromatography: Appearance of new segment of tyrosinases in melanoma cells following interrupted melanogenesis induced by glycosylation inhibitors. *J. Invest. Dermatol.* 85: 165-168, 1985.
- 1-5. Imokawa G., Kawai M, Motegi I, Mishima Y.: Differential analysis of experimental hypermelanosis induced by UVB PUVA and allergic contact dermatitis using brownish guinea pig model. *Arch. Dermatol. Res.* 278: 352-362, 1986.
- 1-6. Imokawa G., Mishima Y.: Importance of glycoproteins in the: initiation of melanogenesis: An electron microscopic study of B-16 melanoma cells after release from inhibition of glycosylation. *J. Invest. Dermatol.* 87: 319-324, 1986.
- 1-7. Imokawa G., Kawai M.: Differential hypermelanosis induced by allergic contact dermatitis. *J. Invest. Dermatol.* 89: 540-546, 1987.
- 1-8. Imokawa G., Yada Y., Hori Y.: Induction of melanization within hair bulb melanocytes in chinchilla mutant by melanogenic stimulants. *J. Invest. Dermatol.* 91: 106-113, 1988.
- 1-9. Imokawa G.: Analysis of initial melanogenesis including tyrosinase transfer and melanosome differentiation through interrupted melanization by glutathione. *J. Invest. Dermatol.* 93: 100-107, 1989.
- 1-10. Genji Imokawa, Toru Tejima: A possible role of prostaglandins in PUVA-induced inflammation: Implication by organ cultured skin. *J Invest Dermatol* 94: 296-300, 1989.
- 1-11. Imokawa G.: Analysis of carbohydrate properties essential for melanogenesis in

- tyrosinases of cultured melanoma cells by differential carbohydrate processing inhibitions. *J. Invest. Dermatol.* 95: 39-49, 1990
- 1-12. Yada Y., Higuchi K., Imokawa G.: Effects of endothelins on signal transduction and proliferation in human melanocytes. *J. Biol. Chem.* 266: 18352-18357, 1991
 - 1-13. Imokawa G., Yada Y., Okuda M.: Allergic contact dermatitis releases soluble factors that stimulate melanogenesis through activation of protein kinase C-related signal transduction pathway. *J. Invest. Dermatol.* 99: 482-488, 1992.
 - 1-14. Imokawa G., Yada Y., Miyagishi M.: Endothelins secreted from human keratinocytes are intrinsic mitogens for human melanocytes. *J. Biol.Chem.* 267: 24675-24680, 1992.
 - 1-15. Tsukamoto K., Uno A., Kubota Y., Hori Y., Imokawa G.: Role of asparagine-linked carbohydrates in pulmonary metastasis of B16-F10 murine melanoma cells: implication through glycosylation inhibition by nojirimycin. *Melanoma Res.* 2: 33-39, 1992.
 - 1-16. Imokawa G., Motegi I.: Skin organ culture model for examining epidermal melanization. *J. Invest. Dermatol.* 100: 47-54, 1993.
 - 1-17. Kobayashi T., Urabe K., Winder A., Jimenez-Cervantes C., Imokawa G., Brewington T., Solano F., Garcia-Borron JC., Hearing VC.: Tyrosinase related protein 1 (TRP1) function as a DHICA oxidase in melanin biosynthesis. *EMBO Journal* 13: 5818-5825, 1994.
 - 1-18. Kobayashi T, Vieira WD, Potterf B, Sakai C, Imokawa G, Hearing VJ. Modulation of melanogenic protein expression during the switch from eu- to pheomelanogenesis. *J Cell Sci.* 1995; 108: 2301-2309.
 - 1-19. Imokawa G, Miyagishi M, Yada Y: Endothelin-1 as a new melanogen: coordinated expression of its gene and the tyrosinase gene in UVB-exposed human epidermis. *J Invest Dermatol* 105:32-37, 1995
 - 1-20. Imokawa G., Yada Y., Morisaki N., Kimura M.: Granulocyte/macrophage-colony stimulatory factor is an intrinsic keratinocyte-derived growth factor for human melanocytes in UVA-induced melanosis. *Biochem. J.* 313: 625-631, 1996.
 - 1-21. Imokawa G., Yada Y., Kimura M.: Signaling mechanisms of endothelin-induced mitogenesis and melanogenesis in human melanocytes. *Biochem. J.* 314: 305-312, 1996.
 - 1-22. Nakagawa H., Imokawa G.: Characterization of melanogenesis in normal human epidermal melanocytes by chemical and ultrastructural analysis. *Pigment Cell Res.* 9: 175-178, 1996.
 - 1-23. Teraki E., Tajima S., Manaka I., Kawashima M., Miyagishi M., Imokawa G.: Role of endothelin-1 in hyperpigmentation in seborrhoeic keratosis. *Brit. J. Dermatol.* 135:

918-923, 1996.

- 1-24. Imokawa G., Kobayashi T., Miyagishi M., Higashi K., Yada Y.: The role of endothelin-1 in epidermal hyperpigmentation and signaling mechanisms of mitogenesis and melanogenesis. *Pigment Cell Res.* 10: 218-228, 1997.
- 1-25. Imokawa G., Yada Y., Higuchi K.: Purification and characterization of an allergy-induced melanogenic stimulating factor in brownish guinea pig skin. *J. Biol. Chem.* 273: 1605-1612, 1998.
- 1-26. Imokawa G., Yada Y., Morisaki N., Kimura M.: Biological characterization of human fibroblast-derived mitogenic factors for human melanocytes. *Biochem. J.* 330: 1235-1239, 1998.
- 1-27. Imokawa G, Kobayashi T, Miyagishi M: Intracellular signaling mechanisms leading to synergistic effects of endothelin-1 and stem cell factor on proliferation of cultured human melanocytes: Cross-talk via trans-activation of the tyrosine kinase c-kit receptor. *J. Biol. Chem.* 275: 33321-33328, 2000.
- 1-28. Hachiya A, Kobayashi A, Ohuchi A, Takema Y, Imokawa G: The paracrine role of stem cell factor/c-kit signaling in the activation of human melanocytes in ultraviolet B-induced pigmentation. *J Invest Dermatol* 116(4): 578-586, 2001
- 1-29. Kadono S, Manaka I, Kawashima M, Kobayashi T, Imokawa G: The role of the epidermal endothelin cascade in the hyperpigmentation mechanism of lentigo senilis. *J Invest Dermatol* 116(4): 571-577, 2001.
- 1-30. Shishido E, Kadono S, Manaka I, Kawashima M, Imokawa G: The mechanism of epidermal hyperpigmentation in dermatofibroma is associated with stem cell factor and hepatocyte growth factor expression. *J Invest Dermatol* 117(3): 627-633, 2001
- 1-31. Manaka I, Kadono S, Kawashima M, Kobayashi T, Imokawa G.: The mechanism of hyperpigmentation in seborrheic keratosis involves the high expression of endothelin converting enzyme-1 and TNF α , which stimulate secretion of endothelin-1. *Br J Dermatol* 145: 895-903, 2001
- 1-32. Hachiya A, Kobayashi T, Takema Y, Imokawa G: Biochemical characterization of endothelin-converting enzyme-1 α in cultured skin-derived cells and its postulated role in the stimulation of melanogenesis in human epidermis. *J Biol Chem* 277(7): 5395-5403, 2002
- 1-33. Okazaki M, Yoshimura K, Suzuki Y, Uchida G, Kitano Y, Harii K, Imokawa G: The mechanism of epidermal hyperpigmentation in Café-Au-Lait Macules of neurofibromatosis type-1 (von Recklinghausen's Disease) may be associated with dermal fibroblast-derived stem cell factor and hepatocyte growth factor. *Br J Dermatol* 148: 689-697, 2003

- 1-34. Higuchi K, Kawashima M, Ichikawa Y, Imokawa G: Sphingosylphosphorylcholine is a melanogenic stimulator for human melanocytes. *Pigment Cell Res.* 16: 670-678, 2003
- 1-35. Kitamura R, Tsukamoto K, Harada K, Shimizu A, Shimada S, Kobayashi T, Imokawa G. Mechanisms underlying the dysfunction of melanocytes in vitiligo epidermis: Role of SCF/KIT protein interactions and its downstream effector, MITF-M. *J Pathol* 202(4): 463-475, 2004
- 1-36. Hattori H, Kawashima M, Ichikawa Y, Imokawa G. The epidermal stem cell factor is over-expressed in lentigo senilis: Implication for the mechanism of hyperpigmentation. *J Invest Dermatol* 122: 1256-1265, 2004
- 1-37. Sakuraba K, Hayashi N, Kawashima M, Imokawa G: Down-regulated PAR-2 is associated in part with interrupted melanosome transfer in pigmented basal cell epithelioma. *Pigment Cell Res* 4: 371-378, 2004
- 1-38. Hachiya A, Kobayashi A, Yoshida Y, Kitahara T, Takema Y, Imokawa G: Biphasic expression of two paracrine melanogenic cytokines, stem cell factor and endothelin-1, in ultraviolet B-induced human melanogenesis. *Am J Pathol.* 65(6):2099-2109, 2004
- 1-39. Kayo Sato-Jin, Emi K. Nishimura, Eijiro Akasaka, Wade Huber, Hajime Nakano, Arlo Miller, Jinyan Du, Min Wu, Katsumi Hanada, Daisuke Sawamura, David E. Fisher, Genji Imokawa. Epistatic Connections between MITF and Endothelin Signaling in Waardenburg Syndrome and Other Pigmentary Disorders. *FASEB J* 22(4):1155-1168, 2008
- 1-40. Junichi Hasegawa, Yasufumi Goto, Hiroshi Murata, Minoru Takata, Toshiaki Saida, Genji Imokawa: Down-regulated melanogenic paracrine cytokine linkages in hypopigmented palmoplantar skin. *Pigment Cell & Melanoma Research* 21(6): 787-699, 2008
- 1-41. Yuki Mizutani, Nobukazu Hayashi, Makoto Kawashima, Genji Imokawa: A single UVB exposure increases the expression of functional KIT in human melanocytes by upregulating MITF expression through the phosphorylation of p38/CREB. *Arch Dermatol Res* 302(4): 283-294, 2010
- 1-42. Hiroaki Nakajima, Yuki Wakabayashi, Kazumasa Wakamatsu, Genji Imokawa: An extract of *Melia toosendan* attenuates endothelin-1-stimulated pigmentation in human epidermal equivalents through the interruption of PKC activity within melanocytes. *Arch Dermatol Res* 303(4):263-276. 2011
- 1-43. Hiroaki Nakajima, Yuki Wakabayashi, Kazumasa Wakamatsu, Genji Imokawa. An extract of *Withania somnifera* attenuates endothelin-1-stimulated pigmentation in human epidermal equivalents through the interruption of PKC activity within

- melanocytes. *Phytotherapy Res* 25: 1398-1411, 2011
- 1-44. Hiroaki Nakajima, Katsunori Fukazawa, Yuki Wakabayashi, Kazumasa Wakamatsu, Genji Imokawa: *Withania somnifera* extract attenuates stem cell factor-stimulated pigmentation in human epidermal equivalent through interruption of ERK phosphorylation within melanocytes. *J Natural Medicine* 66: 435-646, 2012
 - 1-45. Nakajima H, Fukazawa K, Wakabayashi Y, Wakamatsu K, Senda K, Imokawa G: Abrogating effect of a xanthophyll carotenoid astaxanthin on the stem cell factor-induced stimulation of human epidermal pigmentation *Arch Dermatol Res* 304(10):803-816, 2012
 - 1-46. Wakabayashi, Y., Nakajima H., Imokawa, G. Abrogating effect of N-linked carbohydrate modifiers on the stem cell factor and endothelin-1- stimulated epidermal pigmentation in human epidermal equivalents. *J. Dermatol. Sci.* 69(3):215-28, 2013.
 - 1-47. Takenaka Y, Hoshino Y, Nakajima H, Hayashi N, Kawashima M, Imokawa G.: Paracrine cytokine mechanisms underlying the hyperpigmentation of seborrheic keratosis in covered skin areas. *J Dermatol* 40(7):533-542, 2013 doi: 10.1111/1346-8138.12178.
 - 1-48. Hiroaki Nakajima, Takeshi Nagata, Shihiro Koga, Genji Imokawa: Reduced glutathione disrupts the intracellular trafficking of tyrosinase and tyrosinase-related protein-1 but not dopachrome tautomerase and Pmel17 to melanosomes, which results in the attenuation of melanization. *Arch Dermatol Res* 306(1):37-49, 2014 doi: 10.1007/s00403-013-1376-z.
 - 1-49. Hideki Tagashira, Aki Miyamoto, Sei-ichi Kitamura, Masahito Tsubata, Kazuya Yamaguchi, Kinya Takagaki, Genji Imokawa: UVB stimulates the expression of endothelin B receptor in human melanocytes via a sequential activation of the p38/MSK1/CREB/MITF pathway which can be interrupted by a French maritime pine bark extract through a direct inactivation of MSK1. *PLoS One*. 2015 Jun 1;10(6):e0128678. doi: 10.1371/journal.pone.0128678.
 - 1-50. Takao Niwano, Shuko Terazawa, Hiroaki Nakajima, Yuki Wakabayashi, Genji Imokawa: Astaxanthin and withaferin A block paracrine cytokine interactions between UVB exposed human keratinocytes and human melanocytes via the attenuation of endothelin1 secretion and its downstream intracellular signaling. *Cytokine* 73(2):184-197, 2015 doi: 10.1016/j.cyto.2015.02.006.
 - 1-51. Shuko Terazawa, Hiroaki Nakajima, Katsunori Fukasawa, Genji Imokawa: Withaferin A abolishes the stem cell factor-stimulated pigmentation of human epidermal equivalents by interrupting the autophosphorylation of c-KIT in human melanocytes.

- Arch Dermatol Res 307(1): 73-88,. 2015 DOI: 10.1007/s00403-014-1518-y
- 1-52. Takao Niwano, Shuko Terazawa, Hiroaki Nakajima, Genji Imokawa: The stem cell factor-stimulated melanogenesis in human melanocytes can be abrogated by interrupting the phosphorylation of MSK1: Evidence for involvement of the p38/MSK1/CREB/MITF axis. Arch Dermatol Res 310(3): 187–196, 2018
- 1-53. Takao Niwano, Shuko Terazawa, Yoriko Sato, Takahiro Kato, Hiroaki Nakajima, Genji Imokawa: Glucosamine abrogates the stem cell factor + endothelin-1-induced stimulation of melanogenesis via a deficiency in MITF expression due to the proteolytic degradation of CREB in human melanocytes. Arch Dermatol Res 310(8):625-637, 2018 doi: 10.1007/s00403-018-1850-8.
- 1-54. Yuko Ishikawa, Takao Niwano, Shinichi Hirano, Kayoko Numano, Kazunori Takasima, Genji Imokawa: Whitening effect of L-ascorbate-2-phosphate trisodium salt on solar Lentigos. Arch Dermatol Res 311(3):183-191, 2019

2. 光生物学的研究論文

- 2-1. Imayama S., Nakamura K., Takeuchi M., Hori Y., Takema Y., Sakaino Y., Imokawa G.: Ultraviolet-B irradiation deforms the configuration of elastic fibers during the induction of actinic elastosis in rats. J. Dermatol. Sci. 7: 32-38, 1994.
- 2-2. Imokawa G., Takema Y., Yorimoto Y., Tsukahara K., Kawai M., Imayama S.: Degree of ultraviolet-induced tortuosity of elastic fibers in rat skin is age-dependent. J. Invest. Dermatol. 105: 254-258, 1995.
- 2-3. Takema Y., Fujimura T., Ohsu H., Imokawa G.: Unusual wrinkle formation after temporary skin fixation followed by UVB irradiation in hairless mouse skin. Experimental Dermatology 5: 145-149, 1996.
- 2-4. Takema Y., Imokawa G.: The effects of UV-A and UV-B irradiation on the viscoelastic properties of hairless mouse skin in vivo. Dermatology 196: 397-400, 1998.
- 2-5. Tsukahara K., Takema Y., Fujimura T., Moriwaki S., Kitahara T., Imayama S., Imokawa G.: All-trans retinoic acid promotes the repair of tortuosity of elastic fibers in rat skin. Br. J. Dermatol. 140: 1048-1053, 1999.
- 2-6. Tsukahara K, Takema Y, Yorimoto Y, Moriwaki S, Fujimura T, Imokawa G: Effects of single shot UVB irradiation on skin elasticity and three-dimensional microstructure of elastic fibers in rat ski. Photomedicine and photobiology 21: 77-79, 1999
- 2-7. Tsukahara K., Takema Y., Moriwaki S, Fujimura T, Imayama S., Imokawa G.: Carbon

- Dioxide Laser Treatment Promotes Repair of the Three-Dimensional Network of Elastic Fibers in Rat Skin. *Br J Dermatol* 144(3): 452-458, 2001
- 2-8. Tsukahara K, Takema Y, Moriwaki S, Tsuji N, Suzuki Y, Fujimura T, Imokawa G. Selective inhibition of skin fibroblast elastase elicits a dose- dependent prevention of UVB-induced wrinkle formation. *J Invest Dermatol* 117(3): 671-677, 2001
- 2-9. Tsuji N, Moriwaki S, Suzuki Y, Takema Y, Imokawa G.: The role of elastases secreted by fibroblasts in wrinkle formation: Implication through selective inhibition of elastase activity. *Photochem Photobiol* 74(2):283-290, 2001
- 2-10. Akazaki S, Nakagawa H, Kazama H, Osanai O, Kawai M, Takema Y, Imokawa G: Age-related changes in skin wrinkles assessed by a novel three-dimensional morphometric analysis. *Br J Dermatol* 147: 689-695, 2002
- 2-11. Tsukahara K, Nakagawa H, Moriwaki S, Shingo K, Ohuchi A, Takema Y, Imokawa G: Ovariectomy is sufficient to accelerate spontaneous skin ageing and to stimulate ultraviolet irradiation-induced photoageing of murin skin. *Br J Dermatol*. 151(5):984-994, 2004
- 2-12. Tsukahara K, Nakagawa H, Moriwaki S, Takema Y, Fujimura T, Imokawa G: Inhibition of ultraviolet-B-induced wrinkle formation by an elastase-inhibiting herbal extract: Implication for the mechanism underlying elastase-associated wrinkles. *International J Dermatol*, 45: 460-468, 2004
- 2-13. Fujimura T, Moriwaki S, Imokawa G, Takema Y: Crucial role of fibroblast integrins alpha2 and beta1 in maintaining the structural and mechanical properties of the skin. *J Dermatol Sci* 45(1): 45-53, 2007
- 2-14. Kaoru Suganuma, Hiroaki Nakajima, Mamitaro Otsuki, Genji Imokawa: Astaxanthin attenuates the UVA-induced up-regulation of matrix-metalloproteinase-1 and skin fibroblast elastase in human dermal fibroblasts. *J Dermatol Sci* 58(2): 136-142, 2010
- 2-15. Naoko Morisaki, Shigeru Moriwaki, Yoriko Sugiyama-Nakagiri, Keiichi Haketa, Yoshinori Takema, Genji Imokawa: Nepriylsin is identical to skin fibroblast elastase – its role in skin ageing and UV responses. *J Biol Chem* 285(51): 39819-39827, 2010
- 2-16. Hiroaki Nakajima, Yoshiyuki Ezaki, Tomoyashu Nagai, Ryosuke Yoshioka, Genji Imokawa: Epithelial-mesenchymal interaction during UVB-induced up-regulation of neutral endopeptidase. *Biochem J*. 443(1): 297-305, 2012
- 2-17. Shuko Terazawa, Hiroaki Nakajima, Keisuke Tobita, Genji Imokawa: The decreased secretion of hyaluronan by older human fibroblasts under physiological conditions is mainly associated with the down-regulated expression of hyaluronan synthases but not with the expression levels of hyaluronidases. *Cytotechnology*. 67(4):609-620, 2015 doi: 10.1007/s10616-014-9707-2.

- 2-18. Hiroaki Nakajima, Shuko Terazawa, Takao Niwano, Yorihiro Yamamoto, Genji Imokawa: The Inhibitory Effects of Anti-Oxidants on Ultraviolet-Induced Up-Regulation of the Wrinkling-Inducing Enzyme Neutral Endopeptidase in Human Fibroblasts. PLoS One. 2016 DOI:10.1371/journal.pone.0161580 September 20, 2016
- 2-19. Shuko Terazawa, Mariko Takada, Yoriko Sato, Hiroaki Nakajima, Genji Imokawa: The attenuated secretion of hyaluronan by UVA-exposed human fibroblasts is associated with up- and down-regulation of HYBID and HAS2 expression via activated and inactivated signaling of the p38/ATF2 and JAK2/STAT3 cascades. Int. J. Mol. Sci. 2021 Feb 19;22(4):2057. doi: 10.3390/ijms22042057.