

2. シミメカニズム関連

2021/03/12 現在

ナンバーが赤字表記の論文は送付可能

(請求先：imokawag@dream.ocn.ne.jp)

2-1.シミ関連英語原著論文 (79 報)

1. Imokawa G., Mishima Y.: Biochemical and ultrastructural heterogeneity of so-called Golgi apparatus in the melanin synthesizing cells. Proc. C. Derm. Res. 1: 41- 42, 1976.
2. Imokawa G., Mishima Y.: Inhibitory effect of glucosamine on melanogenesis in vitro. Pro. C. Derm. Res. 4: 33- 34, 1979.
3. Mishima Y., Imokawa G.: Functional and three-dimensional differentiation of smooth membrane structures in melanogenesis. Pigment Cell 4: 277- 290, 1979.
4. Mojamdar M., Imokawa G., Mishima Y.: Electrophoretic characterization of tyrosinase in the coated vesicles of melanoma. J. Dermatol. 6: 5-8, 1979.
5. Imokawa G., Mishima Y.: Isolation and characterization of tyrosinase inhibitors and their differential action on melanogenic subcellular compartments in amelanotic and melanotic melanomas. Brit. J. Dermatol 3: 625-633, 1980.
6. Imokawa G., Mishima Y.: Macromolecular analysis of glucosamine-inhibition in cultured melanoma cells. Pro. Jpn. Invest. Soc. Derm. 5: 103-104, 1980.
7. Imokawa G., Mishima Y.: Isolation and biochemical characterization of tyrosinase-rich GERL and coated vesicle in melanin synthesizing cells. Brit J. Dermatol. 104: 169-178, 1981.
8. Imokawa G., Mishima Y.: Biochemical characterization of tyrosinase inhibitors using tyrosinase binding affinity chromatography. Brit. J. Dermatol. 104: 531-539, 1981.
9. Imokawa G., Mishima Y.: Tyrosinase analysis of glycosylation inhibitor-induced interrupted melanogenesis in cultured melanoma cell. Pro. Jpn. Soc. Invest. Derm. 6: 107-108, 1981.
10. Imokawa G., Mishima Y.: Ultrastructural analysis of interrupted melanogenesis in cultured malignant melanoma cells by glycosylation inhibitors. J. Clin. Elec. Micro. 15: 465-466, 1982.
11. Imokawa G., Mishima Y.: Loss of melanogenic properties in tyrosinases induced by glycosylation inhibitors within malignant melanoma cells. Cancer Res. 42: 1994- 2002, 1982.
12. Mishima Y., Imokawa G.: Selective aberration and pigment loss in melanosomes of malignant melanoma cells in vitro by glycosylation inhibitors: Premelanosomes as glycoprotein. J. Invest. Dermatol. 81: 106-114, 1983.

13. Imokawa G., Mishima Y.: Cellular pharmacology of pigment cells: functional analysis of melanogenic inhibitory agents using synchronized recovery process from depigmentation induced by glycosylation inhibition. Proc. Jpn. Invest. Dermatol. 8: 102-104, 1984.
14. Imokawa G., Mishima Y.: Functional analysis of tyrosinase isozymes of cultured melanoma cells during the recovery period following interrupted melanogenesis induced by glycosylation inhibitors J. Invest. Dermatol. 83:196-201, 1984.
15. Imokawa G., Mishima Y.: Selective alteration of secretory glycoprotein premelanosome and Golgi cisternae induced by carbohydrate processing inhibitor, monensin. Pro. Jpn. Invest. Dermatol. 9: 11-12, 1985.
16. Imokawa G., Mishima Y.: Analysis of tyrosinase as asparagine-linked oligosaccharides by Concanavalin-A lectin chromatography: Appearance of new segment of tyrosinases in melanoma cells following interrupted melanogenesis induced by glycosylation inhibitors. J. Invest. Dermatol. 85: 165-168, 1985.
17. Mishima Y., Imokawa G.: Role of glycosylation in initial melanogenesis: post-inhibition dynamics. Pigment Cell 6: 17-30, 1985.
18. Imokawa G., Kawai M., Motegi I, Mishima Y.: Differential analysis of experimental hypermelanosis induced by UVB PUVA and allergic contact dermatitis using brownish guinea pig model. Arch. Dermatol. Res. 278: 352-362, 1986.
19. Imokawa G., Mishima Y.: Importance of glycoproteins in the: initiation of melanogenesis: An electron microscopic study of B-16 melanoma cells after release from inhibition of glycosylation. J. Invest. Dermatol. 87: 319-324, 1986.
20. Imokawa G., Kawai M.: Differential hypermelanosis induced by allergic contact dermatitis. J. Invest. Dermatol. 89: 540-546, 1987.
21. Imokawa G., Yada Y., Hori Y.: Induction of melanization within hair bulb melanocytes in chinchilla mutant by melanogenic stimulants. Structure and Function of Melanin 4: 60-68, 1987.
22. Imokawa G., Yada Y., Hori Y.: Induction of melanization within hair bulb melanocytes in chinchilla mutant by melanogenic stimulants. Pro. Jap. J. Invest. Dermatol. 11: 91- 92, 1987.
23. Imokawa G., Yada Y., Hori Y.: Induction of melanization within hair bulb melanocytes in chinchilla mutant by melanogenic stimulants. J. Invest. Dermatol. 91: 106-113, 1988.
24. Imokawa G.: Analysis of initial melanogenesis including tyrosinase transfer and melanosome differentiation through interrupted melanization by glutathione. J. Invest. Dermatol. 93: 100-107, 1989.
25. Genji Imokawa, Toru Tejima: A possible role of prostaglandins in PUVA-induced inflammation: Implication by organ cultured skin. J Invest Dermatol 94: 296-300, 1989.

26. Imokawa G.: Analysis of carbohydrate properties essential for melanogenesis in tyrosinases of cultured melanoma cells by differential carbohydrate processing inhibitions. *J. Invest. Dermatol.* 95: 39-49, 1990
27. Yada Y., Higuchi K., Imokawa G.: Effects of endothelins on signal transduction and proliferation in human melanocytes. *J. Biol. Chem.* 266: 18352-18357, 1991
28. Imokawa G., Yada Y., Okuda M.: Allergic contact dermatitis releases soluble factors that stimulate melanogenesis through activation of protein kinase C-related signal-transduction pathway. *J. Invest. Dermatol.* 99: 482-488, 1992.
29. Imokawa G., Yada Y., Miyagishi M.: Endothelins secreted from human keratinocytes are intrinsic mitogens for human melanocytes. *J. Biol.Chem.* 267: 24675-24680, 1992.
30. Tsukamoto K., Uno A., Kubota Y., Hori Y., Imokawa G.: Role of asparagine-linked carbohydrates in pulmonary metastasis of B16-F10 murine melanoma cells: implication through glycosylation inhibition by nojirimycin. *Melanoma Res.* 2: 33-39, 1992.
31. Imokawa G., Motegi I.: Skin organ culture model for examining epidermal melanization. *J. Invest. Dermatol.* 100: 47-54, 1993.
32. Komiyama K., Takamatsu S., Takahashi Y., Shinose M., Hayashi M., Tanaka H., Iwai Y., Omura S., Imokawa G.: New inhibitors of melanogenesis, OH-3984 K1 and K2, I. Taxonomy, fermentation, isolation and biological characteristics. *J. Antibiotics* 46: 1520-1525, 1993.
33. Takamatsu S., Rho MC., Hayashi M., Komiyama K., Tanaka H., Omura S., Imokawa G.: New inhibitor of melanogenesis, OH-3984 K1 and K2, II. Physico-chemical properties and structural elucidation. *J. Antibiotics* 46: 1526-1529, 1993.
34. Stender I.M., Etoj T., Byers H.R., Imokawa G., Tsuchida T., Nakagawa H., Ishibashi Y.: Metastatic behavior and tumorigenicity of a human melanoma cell line (MM-RU) after injection into nude mice. *J. Dermatology (Tokyo)* 20: 611-617, 1993.
35. Kobayashi T., Urabe K., Winder A., Tsukamoto K., Brewington T., Imokawa G., Potterf B., Hearing VJ.: The DHICA oxidase activity of TRP1 and interactions with other melanogenic enzymes. *Pigment Cell Res.*7: 227-234, 1994.
36. Kobayashi T., Urabe K., Winder A., Jimenez-Cervantes C., Imokawa G., Brewington T., Solano F., Garcia-Borron JC., Hearing VC.: Tyrosinase related protein 1 (TRP1) function as a DHICA oxidase in melanin biosynthesis. *EMBO Journal* 13: 5818-5825, 1994.
37. Kobayashi T., Urabe K., Orlow S.J., Higashi K., Imokawa G., Kwon BS, Potterf B., Hearing VJ.: The Pmel 17/Silver Locus Protein, Characterization and investigation of its melanogenic function. *J. Biol. Chem.* 269: 29198-29205, 1994.
38. Kobayashi T, Vieira WD, Potterf B, Sakai C, Imokawa G, Hearing VJ. Modulation of melanogenic protein expression during the switch from eu- to

- pheomelanogenesis. *J Cell Sci.* 1995; 108: 2301-2309.
39. Imokawa G, Miyagishi M, Yada Y: Endothelin-1 as a new melanogen: coordinated expression of its gene and the tyrosinase gene in UVB-exposed human epidermis. *J Invest Dermatol* 105:32-37, 1995
 40. Imokawa G., Yada Y., Morisaki N., Kimura M.: Granulocyte/macrophage-colony-stimulatory factor is an intrinsic keratinocyte-derived growth factor for human melanocytes in UVA-induced melanosis. *Biochem. J.* 313: 625-631, 1996.
 41. Imokawa G., Yada Y., Kimura M.: Signaling mechanisms of endothelin-induced mitogenesis and melanogenesis in human melanocytes. *Biochem. J.* 314: 305-312, 1996.
 42. Takamatsu S., Kim Y.P., Hayashi M., Komiyama K., Imokawa G., Omura S.: A new inhibitor of melanogenesis, albocycline K3, produced by *Streptomyces* sp. OH-3984. *J Antibiot. (Tokyo)* 49: 485-486, 1996.
 43. Nakagawa H., Imokawa G.: Characterization of melanogenesis in normal human epidermal melanocytes by chemical and ultrastructural analysis. *Pigment Cell Res.* 9: 175-178, 1996.
 44. Teraki E., Tajima S., Manaka I., Kawashima M., Miyagishi M., Imokawa G.: Role of endothelin-1 in hyperpigmentation in seborrheic keratosis. *Brit. J. Dermatol.* 135: 918-923, 1996.
 45. Imokawa G., Kobayashi T., Miyagishi M., Higashi K., Yada Y.: The role of endothelin-1 in epidermal hyperpigmentation and signaling mechanisms of mitogenesis and melanogenesis. *Pigment Cell Res.* 10: 218-228, 1997.
 46. Sakai C., Ollmann M., Kobayashi T., Muller J., Vieira D., Imokawa G., Barsh G.S., Hearing V.J.: Modulation of murine melanocyte function in vitro by agouti signal protein. *EMBO J.* 16: 3544-3552, 1997.
 47. Imokawa G., Yada Y., Higuchi K.: Purification and characterization of an allergy-induced melanogenic stimulating factor in brownish guinea pig skin. *J. Biol. Chem.* 273: 1605-1612, 1998.
 48. Imokawa G., Yada Y., Morisaki N., Kimura M.: Biological characterization of human fibroblast-derived mitogenic factors for human melanocytes. *Biochem. J.* 330: 1235-1239, 1998.
 49. Kabayashi T., Imokawa G., Bennett D.C., Hearing V.J.: Tyrosinase stabilization by Tyrp1 Brown locus (the brown locus protein). *J. Biol. Chem.* 273: 31801-31805, 1998.
 50. Imokawa G, Kobayashi T, Miyagishi M: Intracellular signaling mechanisms leading to synergistic effects of endothelin-1 and stem cell factor on proliferation of cultured human melanocytes: Cross-talk via trans-activation of the tyrosine kinase c-kit receptor. *J. Biol. Chem.* 275: 33321-33328, 2000.

51. Hachiya A, Kobayashi A, Ohuchi A, Takema Y, Imokawa G: The paracrine role of stem cell factor/c-kit signaling in the activation of human melanocytes in ultraviolet B-induced pigmentation. *J Invest Dermatol* 116(4): 578-586, 2001
52. Kadono S, Manaka I, Kawashima M, Kobayashi T, Imokawa G: The role of the epidermal endothelin cascade in the hyperpigmentation mechanism of lentigo senilis. *J Invest Dermatol* 116(4): 571-577 2001
53. Shishido E, Kadono S, Manaka I, Kawashima M, Imokawa G: The mechanism of epidermal hyperpigmentation in dermatofibroma is associated with stem cell factor and hepatocyte growth factor expression. *J Invest Dermatol* 117(3), 627-633, 2001
54. Manaka I, Kadono S, Kawashima M, Kobayashi T, Imokawa G.: The mechanism of hyperpigmentation in seborrhoeic keratosis involves the high expression of endothelin-converting enzyme-1 α and TNF α , which stimulate secretion of endothelin-1. *Br J Dermatol* 145: 895-903, 2001
55. Hachiya A, Kobayashi T, Takema Y, Imokawa G: Biochemical characterization of endothelin-converting enzyme-1 α in cultured skin-derived cells and its postulated role in the stimulation of melanogenesis in human epidermis. *J Biol Chem* 277(7): 5395-5403, 2002
56. Okazaki M, Yoshimura K, Suzuki Y, Uchida G, Kitano Y, Harii K, Imokawa G: The mechanism of epidermal hyperpigmentation in Café-Au-Lait Macules of neurofibromatosis type-1 (von Recklinghausen's Disease) may be associated with dermal fibroblast-derived stem cell factor and hepatocyte growth factor. *Br J Dermatol* 148: 689-697, 2003
57. Higuchi K, Kawashima M, Ichikawa Y, Imokawa G: Sphingosylphosphorylcholine is a melanogenic stimulator for human melanocytes. *Pigment Cell Res.* 16: 670-678., 2003
58. Kitamura R, Tsukamoto K, Harada K, Shimizu A, Shimada S, Kobayashi T, Imokawa G. Mechanisms underlying the dysfunction of melanocytes in vitiligo epidermis: Role of SCF/KIT protein interactions and its downstream effector, MITF-M. *J Pathol* 202(4): 463-475, 2004
59. Hattori H, Kawashima M, Ichikawa Y, Imokawa G. The epidermal stem cell factor is over-expressed in lentigo senilis: Implication for the mechanism of hyperpigmentation. *J Invest Dermatol* 122: 1256-1265, 2004
60. Sakuraba K, Hayashi N, Kawashima M, Imokawa G: Down-regulated PAR-2 is associated in part with interrupted melanosome transfer in pigmented basal cell epithelioma. *Pigment Cell Res* 4: 371-378, 2004
61. Hachiya A, Kobayashi A, Yoshida Y, Kitahara T, Takema Y, Imokawa G: Biphasic expression of two paracrine melanogenic cytokines, stem cell factor and endothelin-1, in

- ultraviolet B-induced human melanogenesis. *Am J Pathol.* 2004 Dec;1 65(6):2099-2109. 2004
62. Kayo Sato-Jin, Emi K. Nishimura, Eijiro Akasaka, Wade Huber, Hajime Nakano, Arlo Miller, Jinyan Du, Min Wu, Katsumi Hanada, Daisuke Sawamura, David E. Fisher, **Genji Imokawa**. Epistatic Connections between MITF and Endothelin Signaling in Waardenburg Syndrome and Other Pigmentary Disorders. *FASEB J* 22(4):1155-68. 2008
 63. Junichi Hasegawa, Yasufumi Goto, Hiroshi Murata, Minoru Takata, Toshiaki Saida, **Genji Imokawa**: Down-regulated melanogenic paracrine cytokine linkages in hypopigmented palmoplantar skin. *Pigment Cell & Melanoma Research* 21(6) 787-699. 2008
 64. Yuki Mizutani, Nobukazu Hayashi, Makoto Kawashima, **Genji Imokawa**: A single UVB exposure increases the expression of functional KIT in human melanocytes by up-regulating MITF expression through the phosphorylation of p38/CREB. *Arch Dermatol Res* 2010 May;302(4):283-94.
 65. Hiroaki Nakajima, Yuki Wakabayashi, Kazumasa Wakamatsu, **Genji Imokawa**: An extract of *Melia toosendan* attenuates endothelin-1-stimulated pigmentation in human epidermal equivalents through the interruption of PKC activity within melanocytes. *Arch Dermatol Res* (2011) 303:263–276147.
 66. Hiroaki Nakajima, Yuki Wakabayashi, Kazumasa Wakamatsu, **Genji Imokawa**. An extract of *Withania somnifera* attenuates endothelin-1-stimulated pigmentation in human epidermal equivalents through the interruption of PKC activity within melanocytes. *Phytotherapy Res.* 2011; 25: 1398-411.
 67. Hiroaki Nakajima, Katsunori Fukazawa, Yuki Wakabayashi, Kazumasa Wakamatsu, **Genji Imokawa**: *Withania somnifera* extract attenuates stem cell factor-stimulated pigmentation in human epidermal equivalent through interruption of ERK phosphorylation within melanocytes. *J Natural Medicine* 2012; 66: 435-46.
 68. Nakajima H, Fukazawa K, Wakabayashi Y, Wakamatsu K, Senda K, **Imokawa G**: Abrogating effect of a xanthophyll carotenoid astaxanthin on the stem cell factor-induced stimulation of human epidermal pigmentation *Arch Dermatol Res* 2012 Dec;304(10):803-16.
 69. Wakabayashi, Y., Nakajima H., **Imokawa, G**. Abrogating effect of N-linked carbohydrate modifiers on the stem cell factor and endothelin-1- stimulated epidermal pigmentation in human epidermal equivalents. *J. Dermatol. Sci.* 69(3):215-28, 2013.
 70. Takenaka Y, Hoshino Y, Nakajima H, Hayashi N, Kawashima M, **Imokawa G**.: Paracrine cytokine mechanisms underlying the hyperpigmentation of seborrheic keratosis in covered skin areas. *J Dermatol.* 2013; 40(7):533-42. doi: 10.1111/1346-8138.12178.

71. Bong-Keun Choi, Byung-Yoon Cha, Takeshi Fujiwara, Akihiko Kanamoto, Je-Tae Woo, Makoto Ojika, **Genji Imokawa**: Arenarol isolated from a marine sponge abrogates endothelin-1-stimulated melanogenesis by interrupting MEK phosphorylation in normal human melanocytes. *Cytotechnology* 2013; 65(6):915-26.
72. Hiroaki Nakajima, Takeshi Nagata, Shihiro Koga, **Genji Imokawa**: Reduced glutathione disrupts the intracellular trafficking of tyrosinase and tyrosinase-related protein-1 but not dopachrome tautomerase and Pmel17 to melanosomes, which results in the attenuation of melanization. *Arch Dermatol Res* 2014 Jan;306(1):37-49. doi: 10.1007/s00403-013-1376-z.
73. Hideki Tagashira, Aki Miyamoto, Sei-ichi Kitamura, Masahito Tsubata, Kazuya Yamaguchi, Kinya Takagaki, **Genji Imokawa**: UVB stimulates the expression of endothelin B receptor in human melanocytes via a sequential activation of the p38/MSK1/CREB/MITF pathway which can be interrupted by a French maritime pine bark extract through a direct inactivation of MSK1. *PLoS One*. 2015 Jun 1;10(6):e0128678. doi: 10.1371/journal.pone.0128678.
74. Takao Niwano, Shuko Terazawa, Hiroaki Nakajima, Yuki Wakabayashi, **Genji Imokawa**: Astaxanthin and withaferin A block paracrine cytokine interactions between UVB-exposed human keratinocytes and human melanocytes via the attenuation of endothelin-1 secretion and its downstream intracellular signaling. *Cytokine*. 2015 Mar 13;73(2):184-197. doi: 10.1016/j.cyto.2015.02.006.
75. Shuko Terazawa, Hiroaki Nakajima, Katsunori Fukasawa, **Genji Imokawa**: Withaferin A abolishes the stem cell factor-stimulated pigmentation of human epidermal equivalents by interrupting the autophosphorylation of c-KIT in human melanocytes. *Arch Dermatol Res* 2015; 307(1) 73-88. DOI: 10.1007/s00403-014-1518-y
76. Han C, Furukawa H, Tomura T, Fudou R, Kaida K, Choi BK, **Imokawa G**, Ojika M. Bioactive Maleic Anhydrides and Related Diacids from the Aquatic Hyphomycete *Tricladium castaneicola*. *J Nat Prod*. 2015 Apr 24;78(4):639-44. doi: 10.1021/np500773s.
77. Takao Niwano, Shuko Terazawa, Hiroaki Nakajima, **Genji Imokawa**: The stem cell factor-stimulated melanogenesis in human melanocytes can be abrogated by interrupting the phosphorylation of MSK1: Evidence for involvement of the p38/MSK1/CREB/MITF axis. *Arch Dermatol Res* 2018; 310(3), 187–196.
78. Takao Niwano, Shuko Terazawa, Yoriko Sato, Takahiro Kato, Hiroaki Nakajima, **Genji Imokawa**: Glucosamine abrogates the stem cell factor + endothelin-1-induced stimulation of melanogenesis via a deficiency in MITF expression due to the proteolytic degradation of CREB in human melanocytes. *Arch Dermatol Res*. 2018

Oct;310(8):625-637. doi: 10.1007/s00403-018-1850-8.

79. Yuko Ishikawa, Takao Niwano, Shinichi Hirano, Kayoko Numano, Kazunori Takasima, **Genji Imokawa**: Whitening effect of L-ascorbate-2-phosphate trisodium salt on solar lentigos. Arch Dermatol Res. 2019; 311(3):183-191.

2-2.シミ関連日本語原著論文 (3報)

1. 市橋正光、小林明美、奥田峰広、**芋川玄爾**: カミツレエキスの紫外線誘導色素沈着に対する抑制効果。皮膚 41(4): 475-480, 1999
2. 川島 眞、奥田峰広、小林明美、**芋川玄爾**、紫外線誘導色素沈着に対するカミツレエキスの抑制効果。西日本皮膚科 61(5) 682-685, 1999.
3. 赤坂 英二郎、中野 創、神 可代、木村 一之、六戸 大樹、**芋川玄爾**、澤村 大輔: 紫外線防御機構におけるメラノサイト, ケラチノサイトの エンドセリン-1/幹細胞因子を介した相互作用 弘前医学 62:138-143, 2011

2-3.シミ関連英語総説論文 (10報)

1. **Imokawa G**: Autocrine and paracrine regulation of melanocytes in human skin and in pigmentary disorders. Pigment Cell Res 17: 96-110, 2004
2. **Imokawa, G.**; Ishida, K. Inhibitors of Intracellular Signaling Pathways that Lead to Stimulated Epidermal Pigmentation: Perspective of Anti-Pigmenting Agents. Int. J. Mol. Sci. 2014, 15, 8293-8315. doi:10.3390/ijms15058293
3. Shuko Terazawa, **Genji Imokawa**: Signaling cascades activated by UVB in human melanocytes lead to the increased expression of melanocyte-receptors, endothelin B receptor and c-KIT. Photochem Photobiol. 2017 Oct 4. doi: 10.1111/php.12848.
4. **Genji Imokawa**: Mechanism of melanocyte activation in seborrheic keratosis. S Geriatr Dermatol Res 2017; 12(2): 1-14
5. Wu S, **Imokawa G**. To be or not to be photopigmented, that is the question: an introduction to the special issue dedicated to photopigmentation and melanoma. Photochem Photobiol. 2018 May;94(3):407-408. Doi:10.1111/php. 12933.
6. **Genji Imokawa**: The xanthophyll carotenoid astaxanthin has distinct biological effects to prevent the photo-aging of the skin even by its post-irradiation treatment. Photochem Photobiol. 2019 Mar;95(2):480-489. doi: 10.1111/php.13034.
7. **Genji Imokawa**: Intracellular signaling mechanisms involved in the biological effects of

the xanthophyll carotenoid astaxanthin to prevent the photo-aging of the skin in a reactive oxygen species depletion-independent manner: The key role of mitogen and stress-activated protein kinase 1" *Photochem Photobiol.* 2019 Mar;95(2):490-500. doi: 10.1111/php.13039.

8. **Genji Imokawa**: The xanthophyll carotenoid astaxanthin has distinct biological effects to prevent ultraviolet light-induced cutaneous inflammation, abnormal keratinization, pigmentation and wrinkling in a reactive oxygen species-depletion independent fashion even by its post-irradiation treatment. *S Geriat Dermatol Res* 2018; 12(2): 1-14
9. **Genji Imokawa**: Intracellular signaling mechanisms involved in the preventive effects of the xanthophyll carotenoid astaxanthin on ultraviolet light-induced cutaneous inflammation, abnormal keratinization, pigmentation and wrinkling in a reactive oxygen species depletion-independent manner: A key role of mitogen and stress-activated protein kinase 1. *S Geriat Dermatol Res* 2019; 12(2): 1-14
10. **Genji Imokawa**: Melanocyte activation mechanisms and rational therapeutic treatment of sola lentigos. *July Int. J. Mol. Sci.* 2019, 20(15), 3666; doi:10.3390/ijms 20153666 <https://doi.org/10.3390/ijms20153666>

2-4.シミ関連日本語総説論文 (37報)

1. 三島豊, **芋川玄爾**, 玉置昭治, 市橋正光, 中西孝文: メラニン生成細胞に関する最近のトピックス. *日本皮膚科会誌* 91: 1463-1464, 1981.
2. 河合道雄, **芋川玄爾**, 桐井範治: 紫外線の皮膚に対する影響とその対策. *C & T*, 3: 3-6, 1988.
3. 河合道雄, 渋谷ゆう子, 桐井範治, **芋川玄爾**: 吸収型日焼け防止剤. *化粧品科学会誌* 13: 231-235, 1989.
4. **芋川玄爾**, 桐井範治: UVA が関与する皮膚反応の特徴と SPF. *Fragrance J.* 9: 41-54, 1992.
5. 河合道雄, **芋川玄爾**: 紫外線の害とその防御法. *皮膚と美容* 24: 7-17, 1992
6. **芋川玄爾**, 木村光利: 美白剤の作用機序とその将来像. *化粧品科学会誌* 17: 149-155, 1993.
7. **芋川玄爾**: 紫外線の皮膚に対する影響と対策. *名古屋テキスタイル研究会* 12: 12-18, 1993.
8. **芋川玄爾**: メラニン産生抑制効果測定法. *Fragrance J.* 13: 18-33, 1994.
9. **芋川玄爾**: 紫外線色素沈着に本質的に関与するヒトメラノサイト増殖因子. *日本皮膚科学会誌* 104: 1625-1628, 1994.
10. **芋川玄爾**: 紫外線による色素沈着とエンドセリン. *臨床皮膚科* 49(5) 70-75, 1995

11. 芋川玄爾: 紫外線色素沈着におけるメラノサイト活性化メカニズム. *Fragrance J.* 14: 38-48, 1995.
12. 鈴木敏幸、芋川玄爾: これからの美白化粧品. *化粧品科学会誌* 19: 183-189, 1996
13. 芋川玄爾: 美白剤の評価. *日本化粧品科学会誌* 20(1) 17-20, 1996
14. 芋川玄爾: 美白剤. *日本皮膚科学会誌* 108: 1690-1694, 1998.
15. 芋川玄爾: 美白剤研究開発の進歩とその課題. *日本化粧品科学会誌* 22: 312-318, 1998.
16. 芋川玄爾: 美白剤の種類と皮膚への効果メカニズム. *皮膚と化粧品. Monthly Book Derma デルマ*, 全日本病院出版会 2000: 11-24
17. 芋川玄爾: シミの予防法: 将来への細胞レベルからのストラテジー. *日本化粧品科学会誌* 24(1): 29-34, 2000.
18. 川島眞、芋川玄爾: 紫外線色素沈着と老人性色素斑における:色素増強メカニズムとカモミラ E T の抗シミ効果. *Monthly Book Derma デルマ (MB Derma)*, 全日本病院出版会 2005: 98: 43-61
19. 芋川玄爾: 機能性化粧品の展望. *Monthly Book Derma デルマ (MB Derma)*, 全日本病院出版会 2007: 132: 175-189
20. 芋川玄爾: ストレスと色素細胞一日焼け遺伝学を中心として 特集1: ヒトの皮膚・眼の色の遺伝学. *遺伝* Vol. 63 No.5 20-26, 2009
21. 芋川玄爾: 老人性色素斑の病態、特に色素沈着増強メカニズムについて. *美容皮膚科学会誌(Aesthetic Dermatology)* 19: 297-309, 2009
22. 芋川玄爾: シミ・シワのメカニズム. *太陽紫外線防御委員会* 19(1) 59-65, 2009
23. 芋川玄爾: グルコサミンの色素細胞へのメラニン合成抑制メカニズム. *グルコサミン研究* 7: 21-26, 2011
24. 芋川玄爾: メラニンからみた美容皮膚科学 -美白剤の展望- *美容皮膚科学会誌 Aesthetic Dermatology* 24: 17-34, 2014
25. 芋川玄爾: 化粧品白斑発症機序の科学的考察: 職業性白斑及び尋常性白斑との違いと類似性. *Fragrance J.* 9: 41-54, 2014.
26. 芋川玄爾: 美白に対する機能評価. *皮膚科の臨床* 56巻 第11号 186-205, 2014
27. 芋川玄爾: 表皮色素異常症におけるパラクリンサイトカイン相互作用メカニズム 第一章: 表皮細胞由来のサイトカインが原因する表皮色素沈着の発症メカニズム—UVB/UVA 色素沈着、老人性色素斑、老人性疣贅—, *日本美容皮膚科学会雑誌 (Aesthetic Dermatology)* Vol.25, No.3 303-321, 2015
28. 芋川玄爾: 表皮性色素異常症におけるパラクリンサイトカイン相互作用メカニズム 第二章: 真皮線維芽細胞由来のサイトカインが原因する表皮性色素沈着の発症メカニズム: 皮膚線維腫・カフェオレ斑・肝斑. *日本美容皮膚科学会雑誌 (Aesthetic Dermatology)* Vol.25, No.4 123-131, 2015
29. 芋川玄爾: 表皮色素異常症におけるパラクリンサイトカイン相互作用メカニズム

第3章：正常皮膚色（人種差および部位差）コントロール，外来ケミカルおよび内因性脂質またはストレス物質が原因する表皮色素異常症のメカニズムー女子顔面黒皮症，アトピー性皮膚炎での dirty skin，手掌足底皮膚の脱色素，尋常性白斑．．日本美容皮膚科学会雑誌（Aesthetic Dermatology） Vol.26：1～20，2016

30. 芋川玄爾：ヒト皮膚の色素形成にかかわる細胞外シグナル因子とヒトメラノサイト細胞内シグナル経路.第1章エンドセリンとエンドセリン B 受容体の相互作用. *Fragrance J.* 9: 57-64, 2017.
31. 芋川玄爾：ヒト皮膚の色素形成にかかわる細胞外シグナル因子とヒトメラノサイト細胞内シグナル経路.第2章ステムセルファクターと c-KIT 受容体 の相互作用. *Fragrance J.* 10: 2-17, 2017.
32. 芋川玄爾：ヒト皮膚の色素形成にかかわる細胞外シグナル因子とヒトメラノサイト細胞内シグナル経路.第3章 GM-CSF/c-MET, GRO α /IL-8R, SPC/EDGR, α MSH/MC1R, bFGF/ FGFR, HGF/HGFR 間の相互作用. *Fragrance J.* 11: 62-71, 2017.
33. 芋川玄爾：老人性色素斑のメラノサイト活性化メカニズムとその合理的な美白法
1章：老人性色素斑のサイトカイン及び受容体発現プロファイル。 *Fragrance J.* 4: 1-28, 2020.
34. 芋川玄爾：老人性色素斑のメラノサイト活性化メカニズムとその合理的な美白法
2章：老人性色素斑の色素増強促進メカニズム. *Fragrance J.* 5: 1-28, 2020.
35. 芋川玄爾：老人性色素斑のメラノサイト活性化メカニズムとその合理的な美白法
3章：老人性色素斑のメラノサイト活性化サイトカインの発現メカニズム. *Fragrance J.* 6: 1-28, 2020.
36. 芋川玄爾：老人性色素斑のメラノサイト活性化メカニズムとその合理的な美白法
4章：メラノサイト細胞内情報伝達メカニズムのブロック剤のスクリーニングとその老人性色素斑への美白効果. *Fragrance J.* 7: 1-28, 2020.
37. 芋川玄爾：老人性色素斑のメラノサイト活性化メカニズムとその合理的な美白法
5章：チロシナーゼ阻害剤リン酸アスコルビン酸ナトリウム塩の老人性色素斑への美白効果. *Fragrance J.* 8: 1-28, 2020.

2-5.シミ関連英語著書（7報）

1. Mishima Y., Ichihashi M., Hayashibe K., Ueda M., Hatta S., Funasaka Y., Imokawa G.: Control of melanogenesis and melanoma oncogenesis. ed. By Bagnara, J.T. In: *Advance in Pigment Cell Research*, Alan R. Liss Inc. 1988: 127- 141
2. Imokawa G, Yada Y, Morisaki N, et al.: Biological characterization of human fibroblast-derived mitogenic factors for human melanocytes. *Biochem J*, 330 : 1235-1239, 1998.

3. **Imokawa G**: Characterization of keratinocyte-and fibroblast-derived mitogens for human melanocyte- their roles in stimulated cutaneous pigmentation. ed. by Y. Hori, V. J. Hearing and J. Nakayama、 Melanogenesis and Malignant Melanoma、 Elsevier Science B. V. Amsterdam, 1996 : pp 35-48
4. **Imokawa G**: Paracrine interactions of melanocytes in pigmentary disorders. Ed. By James J. Nordlund, Raymond E. Boissy, Vincent J Hearing, Richard A. King, William S. Oetting, Jean-Paul Ortonne. The Pigmentary System, Blackwell Publishing, Malden, Massachusetts, USA.2005, pp421-444
5. **Imokawa G**: Vitiligo Book, Vitiligo and Hypopigmentation, II-2i Cytokines and growth factors-II, Blackwell Publishing, Malden, Massachusetts, USA.2008, pp34- Vitiligo 2010, Pages 269-282 Cytokines and growth factors (Book Chapter)
6. **Imokawa G**: Withania Somnifera Extract/Withaferin A as a Prospective Anti-Pigmenting Agent. Science of Ashwagandha: Preventive and Therapeutic Potential, Ed. By Sunil C. Kaul and Renu Wadhwa. Published 2017 by Springer International Publishing AG, pp 121-146.
7. **Imokawa, G.**: Molecular mechanisms of melanocyte activation in solar lentigos and their rational therapeutic treatments /Chapter 2. Cutting Edge on Molecular Mechanisms of Skin Ageing and its Rational Anti-Aging Treatments and Pathogenesis of Atopic Dermatitis. ed. By G Imokawa, Book Publisher International, Third Floor, 207 Regent Street, London, 2021;pp. 146-148 and pp. 172-173. DOI: 10.9734/bpi/mono/978-93-90516-55-1, DOI: <https://doi.org/10.9734/bpi/mono/978-93-90516-55-1>

2-6.シミ関連日本語著書 (12報)

1. **芋川玄爾**, 三島豊: 悪性黒色腫 (hamster) 疾患モデルハンドブック 1981 : 337- 342
2. **芋川玄爾**:美白ーしみは消せるか? スキンケアーの実際, 皮膚科診療プラクティス, 文光堂., 1999: 42-44
3. **芋川玄爾**: 色素沈着メカニズム. 皮膚のメカニズムの解明と計測・評価法、技術情報協会、1999 : 55-81
4. **芋川玄爾** : 顔面に見られる色素沈着症の原因サイトカインとメラノサイト活性化に至る細胞内情報伝達メカニズム。老化防止、美白、保湿化粧品の開発(development of anti-aging, whitening, moisturizing cosmetics) 監修 鈴木正人、シーエムシーpp 69-113、2001
5. **芋川玄爾** : 表皮性色素沈着症のパラクリンサイトカインメカニズムと細胞内情報伝達機構ー美白剤開発の基礎としてー。シーエムシー 2004、監修 鈴木正人、シーエムシー出版 pp 26-67、2004

6. 堀田光行、芋川玄爾：t-フラバノンの育毛効果。アンチ・エイジングシリーズNo. 1 白髪・脱毛・育毛の実際。株式会社 エヌ・ティー・エス。企画編集有限会社ブッカーズ pp 105-116. 2005.
7. 芋川玄爾：美白剤開発のための評価法。バイオテクノロジーシリーズ機能性化粧品素材開発のための実験法—in vitro/細胞/組織培養— 2007 監修 芋川玄爾 シーエムシー出版 pp 167-171.
8. 芋川玄爾：メラノーマ細胞を用いたメラニン抑制剤の評価法②色素再生モデル。バイオテクノロジーシリーズ機能性化粧品素材開発のための実験法—in vitro/細胞/組織培養— 2007 監修 芋川玄爾 シーエムシー出版 pp 194-206.
9. 芋川玄爾：皮膚組織培養を用いた評価法。バイオテクノロジーシリーズ機能性化粧品素材開発のための実験法—in vitro/細胞/組織培養— 2007 監修 芋川玄爾 シーエムシー出版 pp 207-221.
10. 芋川玄爾：毛胞器官培養を用いたメラニン抑制剤評価法。バイオテクノロジーシリーズ機能性化粧品素材開発のための実験法—in vitro/細胞/組織培養— 2007 監修 芋川玄爾 シーエムシー出版 pp 222-229.
11. 芋川玄爾：単著本 ヒト皮膚測定評価試験マニュアル、美白剤。2007 出版社：技術情報協会
12. 芋川玄爾、肥田時征、山下利春：色素細胞：第9章・色素形成にかかわる細胞外シグナルおよび細胞内シグナル経路。発行所：慶応大学出版社 pp56-62